

Mouse Acetyl Coenzyme A Acetyltransferase 2 (ACAT2) ELISA Kit | ACAT2 elisa kit

Mouse Acetyl Coenzyme A Acetyltransferase 2 (ACAT2) ELISA Kit

CATALOG #
AAA67162

REACTIVITY
Mouse


PRODUCT SYNONYMS

Acetyl Coenzyme A Acetyltransferase 2 (ACAT2); Mouse Acetyl Coenzyme A Acetyltransferase 2 (ACAT2) ELISA Kit; ACAT2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of ACAT2. No significant cross-reactivity or interference between ACAT2 and analogues was observed.
Sequence Length	525
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.047ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ACAT2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ACAT2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ACAT2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ACAT2 in mouse tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ACAT2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ACAT2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ACAT2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ACAT2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	22122547
NCBI GeneID	223920
NCBI Accession #	NP_666176.1
NCBI GenBank Nucleotide #	NM_146064.1
UniProt Accession #	O88908; Q8R0Y9
Molecular Weight	60,597 Da
NCBI Official Full Name	sterol O-acyltransferase 2
NCBI Official Synonym Full Names	sterol O-acyltransferase 2
NCBI Official Symbol	Soat2
NCBI Official Synonym Symbols	ACAT2; D15Wsu97e
NCBI Protein Information	sterol O-acyltransferase 2
UniProt Protein Name	Sterol O-acyltransferase 2
UniProt Gene Name	Soat2
UniProt Synonym Gene Names	Acact-2; Acat2; ACAT-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/acetyl-coenzyme-a-acetyltransferase-2-acat2/67162>