

Human Mucin 2 (MUC2) ELISA Kit | MUC2 elisa kit

Human Mucin 2 (MUC2) ELISA Kit

CATALOG #
AAA67179

REACTIVITY
Human


PRODUCT SYNONYMS

Mucin 2 (MUC2); Human Mucin 2 (MUC2) ELISA Kit; MLP SMUC Oligomeric Mucus/Gel-Forming Mucin 2; Intestinal/Tracheal; MUC2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of MUC2. No significant cross-reactivity or interference between MUC2 and analogues was observed.
Sequence Length	5179
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.116ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level MUC2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level MUC2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MUC2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of MUC2 in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MUC2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MUC2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MUC2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of MUC2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	2506877
NCBI GeneID	4583
UniProt Accession #	Q02817; Q14878
Molecular Weight	540,300 Da
NCBI Official Full Name	Mucin-2
NCBI Official Synonym Full Names	mucin 2, oligomeric mucus/gel-forming
NCBI Official Symbol	MUC2
NCBI Official Synonym Symbols	MLP; SMUC; MUC-2
NCBI Protein Information	mucin-2
UniProt Protein Name	Mucin-2
UniProt Gene Name	MUC2
UniProt Synonym Gene Names	SMUC; MUC-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.