

Human ER Lipid Raft Associated Protein 2 (ERLIN2) ELISA Kit | ERLIN2 elisa kit

Human ER Lipid Raft Associated Protein 2 (ERLIN2) ELISA Kit

CATALOG #
AAA67200

REACTIVITY
Human


PRODUCT SYNONYMS

ER Lipid Raft Associated Protein 2 (ERLIN2); Human ER Lipid Raft Associated Protein 2 (ERLIN2) ELISA Kit; Erlin-2; SPFH2; NET32; C8orf2; ER Lipid Raft Associated Protein 2; Stomatin-prohibitin-flotillin-HflC/K domain-containing protein 2; ERLIN2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ERLIN2. No significant cross-reactivity or interference between ERLIN2 and analogues was observed.
Sequence Length	152
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.115ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ERLIN2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ERLIN2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ERLIN2 elisa kit

Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ERLIN2 in human tissue homogenates and other biological fluids.

Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ERLIN2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ERLIN2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ERLIN2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ERLIN2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	51242968
NCBI GeneID	11160
NCBI Accession #	NP_001003790.1
NCBI GenBank Nucleotide #	NM_001003790.3
UniProt Accession #	O94905; Q6NW21; Q86VS6; Q86W49; A0JLQ1; A8K5S9; B4DM38; D3DSW0
Molecular Weight	22,939 Da
NCBI Official Full Name	erlin-2 isoform 2
NCBI Official Synonym Full Names	ER lipid raft associated 2
NCBI Official Symbol	ERLIN2
NCBI Official Synonym Symbols	NET32; SPFH2; SPG18; C8orf2; Erlin-2
NCBI Protein Information	erlin-2
UniProt Protein Name	Erlin-2
UniProt Gene Name	ERLIN2
UniProt Synonym Gene Names	C8orf2; SPFH2; SPFH domain-containing protein 2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/er-lipid-raft-associated-protein-2-erlin2/67200>