

Human Coagulation Factor XII (F12) ELISA Kit | F12 elisa kit

Human Coagulation Factor XII (F12) ELISA Kit

CATALOG #
AAA67330

REACTIVITY
Human


PRODUCT SYNONYMS

Coagulation Factor XII (F12); Human Coagulation Factor XII (F12) ELISA Kit; FXII; HAF; Contact Factor; Hageman Factor; F12 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of F12. No significant cross-reactivity or interference between F12 and analogues was observed.
Sequence Length	615
Samples	Plasma.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.065ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level F12 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level F12 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for F12 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of F12 in human plasma. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to F12. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to F12. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain F12, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of F12 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	145275213
NCBI GeneID	2161
NCBI Accession #	NP_000496.2
NCBI GenBank Nucleotide #	NM_000505.3
UniProt Accession #	P00748; P78339
Molecular Weight	67,792 Da
NCBI Official Full Name	coagulation factor XII preproprotein
NCBI Official Synonym Full Names	coagulation factor XII
NCBI Official Symbol	F12
NCBI Official Synonym Symbols	HAF; HAE3; HAEX
NCBI Protein Information	coagulation factor XII
UniProt Protein Name	Coagulation factor XII
UniProt Gene Name	F12
UniProt Synonym Gene Names	HAF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/coagulation-factor-xii-f12/67330>