

Porcine Hepcidin (Hepc) ELISA Kit | Hepc elisa kit

Porcine Hepcidin (Hepc) ELISA Kit

CATALOG #
AAA67386

REACTIVITY
Porcine


PRODUCT SYNONYMS

Hepcidin (Hepc); Porcine Hepcidin (Hepc) ELISA Kit; HAMP HFE2B PLTR LEAP1 Hepcidin Antimicrobial Peptide Liver-expressed antimicrobial peptide 1 Putative liver tumor regressor; Hepc elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Specificity	This assay has high sensitivity and excellent specificity for detection of Hepc. No significant cross-reactivity or interference between Hepc and analogues was observed.
Sequence Length	82
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	0.31ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Hepc were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Hepc were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Hepc elisa kit	Intended Uses: This kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of Hepc in porcine serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to Hepc. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to Hepc. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Hepc, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of Hepc in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	47523042
NCBI GeneID	397207
NCBI Accession #	NP_999282.1
NCBI GenBank Nucleotide #	NM_214117.1
UniProt Accession #	Q8MJ80
Molecular Weight	8,771 Da
NCBI Official Full Name	hepcidin
NCBI Official Symbol	HAMP
NCBI Protein Information	hepcidin
UniProt Protein Name	Hepcidin
UniProt Gene Name	HAMP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.