

Mouse Interferon Gamma Induced Protein 10kDa (IP10) ELISA Kit | IP10 elisa kit

Mouse Interferon Gamma Induced Protein 10kDa (IP10) ELISA Kit

CATALOG #
AAA67433

REACTIVITY
Mouse


PRODUCT SYNONYMS

Interferon Gamma Induced Protein 10kDa (IP10); Mouse Interferon Gamma Induced Protein 10kDa (IP10) ELISA Kit; CXCL10; C7; IFI10; INP10; SCYB10; Crg2; GIP10; Mob1; Chemokine C-X-C Motif Ligand 10; Small Inducible Cytokine Subfamily B(Cys-X-Cys)Member 10; IP10 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of IP10. No significant cross-reactivity or interference between IP10 and analogues was observed.
Sequence Length	98
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	5.7pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level IP10 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level IP10 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IP10 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of IP10 in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to IP10. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to IP10. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain IP10, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of IP10 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	10946576
NCBI GeneID	15945
NCBI Accession #	NP_067249.1
NCBI GenBank Nucleotide #	NM_021274.2
UniProt Accession #	P17515
Molecular Weight	10,789 Da
NCBI Official Full Name	C-X-C motif chemokine 10
NCBI Official Synonym Full Names	chemokine (C-X-C motif) ligand 10
NCBI Official Symbol	Cxcl10
NCBI Official Synonym Symbols	C7; IP10; CRG-2; INP10; IP-10; Ifi10; mob-1; Scyb10; gIP-10
NCBI Protein Information	C-X-C motif chemokine 10
UniProt Protein Name	C-X-C motif chemokine 10
UniProt Gene Name	Cxcl10
UniProt Synonym Gene Names	Crg2; Ifi10; Inp10; Scyb10; Gamma-IP10; IP-10

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/interferon-gamma-induced-protein-10kda-ip10/67433>