

Human Nerve Growth Factor (NGF) ELISA Kit | NGF elisa kit

Human Nerve Growth Factor (NGF) ELISA Kit

CATALOG #
AAA67471

REACTIVITY
Human


PRODUCT SYNONYMS

Nerve Growth Factor (NGF); Human Nerve Growth Factor (NGF) ELISA Kit; NGFB; Beta-NGF; HSN5; NGF-B; Beta-Nerve Growth Factor; NGF elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of NGF. No significant cross-reactivity or interference between NGF and analogues was observed.
Sequence Length	241
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.1pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level NGF were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level NGF were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NGF elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of NGF in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to NGF. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to NGF. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain NGF, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of NGF in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	70995319
NCBI GeneID	4803
NCBI Accession #	NP_002497.2
NCBI GenBank Nucleotide #	NM_002506.2
UniProt Accession #	P01138; Q6FHA0; Q96P60; Q9P2Q8; Q9UKL8; A1A4E5
Molecular Weight	26,959 Da
NCBI Official Full Name	beta-nerve growth factor
NCBI Official Synonym Full Names	nerve growth factor
NCBI Official Symbol	NGF
NCBI Official Synonym Symbols	NGFB; HSAN5; Beta-NGF
NCBI Protein Information	beta-nerve growth factor
UniProt Protein Name	Beta-nerve growth factor
UniProt Gene Name	NGF
UniProt Synonym Gene Names	NGFB; Beta-NGF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.