

Human Connexin 37 (CX37) ELISA Kit | CX37 elisa kit

Human Connexin 37 (CX37) ELISA Kit

CATALOG #
AAA67622

REACTIVITY
Human


PRODUCT SYNONYMS

Connexin 37 (CX37); Human Connexin 37 (CX37) ELISA Kit; CX37 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of CX37. No significant cross-reactivity or interference between CX37 and analogues was observed.
Sequence Length	333
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.057ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level CX37 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level CX37 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CX37 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CX37 in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CX37. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CX37. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CX37, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of CX37 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	62865885
NCBI GeneID	2701
NCBI Accession #	NP_002051.2
NCBI GenBank Nucleotide #	NM_002060.2
UniProt Accession #	P35212; Q9P106; Q9UBL1; Q9UNA9; Q9UNB0; Q9UNB1; Q9Y5N7; A8K698; D3DPR4
Molecular Weight	37,414 Da
NCBI Official Full Name	gap junction alpha-4 protein
NCBI Official Synonym Full Names	gap junction protein alpha 4
NCBI Official Symbol	GJA4
NCBI Official Synonym Symbols	CX37
NCBI Protein Information	gap junction alpha-4 protein
UniProt Protein Name	Gap junction alpha-4 protein
UniProt Gene Name	GJA4
UniProt Synonym Gene Names	Cx37

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.