

Human Interleukin 17 Receptor A (IL17RA) ELISA Kit | IL17RA elisa kit

Human Interleukin 17 Receptor A (IL17RA) ELISA Kit

CATALOG #
AAA67802

REACTIVITY
Human


PRODUCT SYNONYMS

Interleukin 17 Receptor A (IL17RA); Human Interleukin 17 Receptor A (IL17RA) ELISA Kit; CDw217; CD217; IL17R-A; IL17-RA; IL17R; HIL-17R; IL17RA elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of IL17RA. No significant cross-reactivity or interference between IL17RA and analogues was observed.
Sequence Length	832
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	78.125-5000pg/mL
Sensitivity	30pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level IL17RA were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level IL17RA were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL17RA elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of IL17RA in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to IL17RA. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to IL17RA. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain IL17RA, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of IL17RA in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	583966138
NCBI GeneID	23765
NCBI Accession #	NP_001276834.1
NCBI GenBank Nucleotide #	NM_001289905.1
UniProt Accession #	Q96F46; O43844; Q20WK1
Molecular Weight	92,153 Da
NCBI Official Full Name	interleukin-17 receptor A isoform 2
NCBI Official Synonym Full Names	interleukin 17 receptor A
NCBI Official Symbol	IL17RA
NCBI Official Synonym Symbols	CD217; IL17R; IMD51; CANDF5; CDw217; IL-17RA; hIL-17R
NCBI Protein Information	interleukin-17 receptor A
UniProt Protein Name	Interleukin-17 receptor A
UniProt Gene Name	IL17RA
UniProt Synonym Gene Names	IL17R; IL-17 receptor A; IL-17RA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.