

Human Lymphotoxin Beta Receptor (LTbR) ELISA Kit | LTbR elisa kit

Human Lymphotoxin Beta Receptor (LTbR) ELISA Kit

CATALOG #
AAA67852

REACTIVITY
Human


PRODUCT SYNONYMS

Lymphotoxin Beta Receptor (LTbR); Human Lymphotoxin Beta Receptor (LTbR) ELISA Kit; LTbR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of LTbR. No significant cross-reactivity or interference between LTbR and analogues was observed.
Sequence Length	416
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.104ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level LTbR were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level LTbR were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LTbR elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of LTbR in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to LTbR. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to LTbR. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain LTbR, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of LTbR in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	402534565
NCBI GeneID	4055
NCBI Accession #	NP_001257916.1
NCBI GenBank Nucleotide #	NM_001270987.1
UniProt Accession #	P36941; B7Z1D2; D3DUR2; F5GXE7
Molecular Weight	44,810 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 3 isoform 2
NCBI Official Synonym Full Names	lymphotoxin beta receptor
NCBI Official Symbol	LTBR
NCBI Official Synonym Symbols	CD18; TNFCR; TNFR3; D12S370; TNFR-RP; TNFRSF3; TNFR2-RP; LT-BETA-R; TNF-R-III
NCBI Protein Information	tumor necrosis factor receptor superfamily member 3
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 3
UniProt Gene Name	LTBR
UniProt Synonym Gene Names	D12S370; TNFCR; TNFR3; TNFRSF3; TNF-RIII; TNFR-III

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.