

Human Acrosin (ACR) ELISA Kit | ACR elisa kit

Human Acrosin (ACR) ELISA Kit

CATALOG #
AAA67855

REACTIVITY
Human


PRODUCT SYNONYMS

Acrosin (ACR); Human Acrosin (ACR) ELISA Kit; ACRS; Preproacrosin; ACR elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ACR. No significant cross-reactivity or interference between ACR and analogues was observed.
Sequence Length	421
Samples	Sperm.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.062ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ACR were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ACR were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ACR elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ACR in human sperm. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ACR. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ACR. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ACR, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ACR in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	148613878
NCBI GeneID	49
NCBI Accession #	NP_001088.2
NCBI GenBank Nucleotide #	NM_001097.2
UniProt Accession #	P10323; Q6ICK2
Molecular Weight	45,847 Da
NCBI Official Full Name	acrosin
NCBI Official Synonym Full Names	acrosin
NCBI Official Symbol	ACR
NCBI Protein Information	acrosin
UniProt Protein Name	Acrosin
UniProt Gene Name	ACR
UniProt Synonym Gene Names	ACRS

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/acrosin-acr/67855>