

Mouse Mucin 4 (MUC4) ELISA Kit | MUC4 elisa kit

Mouse Mucin 4 (MUC4) ELISA Kit

CATALOG #
AAA67868

REACTIVITY
Mouse


PRODUCT SYNONYMS

Mucin 4 (MUC4); Mouse Mucin 4 (MUC4) ELISA Kit; ASGP; Mucin 4 Cell Surface Associated; Mucin 4; Tracheobronchial; Testis mucin; Ascites sialoglycoprotein; Pancreatic adenocarcinoma mucin; MUC4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of MUC4. No significant cross-reactivity or interference between MUC4 and analogues was observed.
Sequence Length	3,470
Samples	Tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.087ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level MUC4 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level MUC4 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MUC4 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of MUC4 in mouse tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MUC4. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MUC4. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MUC4, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of MUC4 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	167736365
NCBI GeneID	140474
NCBI Accession #	NP_536705.3
NCBI GenBank Nucleotide #	NM_080457.3
UniProt Accession #	E9Q7Q0; Q8JZM8
Molecular Weight	368,411 Da
NCBI Official Full Name	mucin-4
NCBI Official Synonym Full Names	mucin 4
NCBI Official Symbol	Muc4
NCBI Official Synonym Symbols	Asgp; 4933405I11Rik
NCBI Protein Information	mucin-4
UniProt Protein Name	Mucin-4
UniProt Gene Name	Muc4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.