

Human Relaxin (RLN) ELISA Kit | RLN elisa kit

Human Relaxin (RLN) ELISA Kit

CATALOG #
AAA67930

REACTIVITY
Human


PRODUCT SYNONYMS

Relaxin (RLN); Human Relaxin (RLN) ELISA Kit; RLXH1; Relaxin H1; Prorelaxin H1; RLN elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of RLN. No significant cross-reactivity or interference between RLN and analogues was observed.
Sequence Length	117
Samples	Serum, plasma and other biological fluids.
Assay Type	Competitive inhibition
Detection Range	31.25-2000pg/mL
Sensitivity	9.66pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level RLN were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level RLN were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for RLN elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of RLN in human serum, plasma and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to RLN. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to RLN. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain RLN, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of RLN in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	5902052
NCBI GeneID	6013
NCBI Accession #	NP_008842.1
NCBI GenBank Nucleotide #	NM_006911.3
UniProt Accession #	P04808; Q99936; Q9UQJ1
Molecular Weight	13,011 Da
NCBI Official Full Name	prorelaxin H1 preproprotein
NCBI Official Synonym Full Names	relaxin 1
NCBI Official Symbol	RLN1
NCBI Official Synonym Symbols	H1; H1RLX; RLXH1; bA12D24.3.1; bA12D24.3.2
NCBI Protein Information	prorelaxin H1
UniProt Protein Name	Prorelaxin H1
UniProt Gene Name	RLN1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.