

Human Intercellular Adhesion Molecule 2 (ICAM2) ELISA Kit | ICAM2 elisa kit

Human Intercellular Adhesion Molecule 2 (ICAM2) ELISA Kit

CATALOG #
AAA68040

REACTIVITY
Human


PRODUCT SYNONYMS

Intercellular Adhesion Molecule 2 (ICAM2); Human Intercellular Adhesion Molecule 2 (ICAM2) ELISA Kit; ICAM2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ICAM2. No significant cross-reactivity or interference between ICAM2 and analogues was observed.
Sequence Length	275
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	6.25-400ng/mL
Sensitivity	<1.6ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ICAM2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ICAM2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ICAM2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of ICAM2 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ICAM2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ICAM2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ICAM2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ICAM2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	153082696
NCBI GeneID	3384
NCBI Accession #	NP_000864.2
NCBI GenBank Nucleotide #	NM_000873.3
UniProt Accession #	P13598; Q14600
Molecular Weight	30,654 Da
NCBI Official Full Name	intercellular adhesion molecule 2
NCBI Official Synonym Full Names	intercellular adhesion molecule 2
NCBI Official Symbol	ICAM2
NCBI Official Synonym Symbols	CD102
NCBI Protein Information	intercellular adhesion molecule 2
UniProt Protein Name	Intercellular adhesion molecule 2
UniProt Gene Name	ICAM2
UniProt Synonym Gene Names	ICAM-2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/intercellular-adhesion-molecule-2-icam2/68040>