

Human Lymphocyte Activation Gene 3 (LAG3) ELISA Kit | LAG3 elisa kit

Human Lymphocyte Activation Gene 3 (LAG3) ELISA Kit

CATALOG #
AAA68049

REACTIVITY
Human


PRODUCT SYNONYMS

Lymphocyte Activation Gene 3 (LAG3); Human Lymphocyte Activation Gene 3 (LAG3) ELISA Kit; LAG3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of LAG3. No significant cross-reactivity or interference between LAG3 and analogues was observed.
Sequence Length	360
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.063ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level LAG3 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level LAG3 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for LAG3 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of LAG3 in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to LAG3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to LAG3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain LAG3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of LAG3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	167614500
NCBI GeneID	3902
NCBI Accession #	NP_002277.4
NCBI GenBank Nucleotide #	NM_002286.5
UniProt Accession #	P18627; Q7Z643; A8K7T9
Molecular Weight	39,103 Da
NCBI Official Full Name	lymphocyte activation gene 3 protein
NCBI Official Synonym Full Names	lymphocyte activating 3
NCBI Official Symbol	LAG3
NCBI Official Synonym Symbols	CD223
NCBI Protein Information	lymphocyte activation gene 3 protein
UniProt Protein Name	Lymphocyte activation gene 3 protein
UniProt Gene Name	LAG3
UniProt Synonym Gene Names	FDC; LAG-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-elisa-kits/lymphocyte-activation-gene-3-lag3/68049>