

Human Rh Blood Group, D Antigen (RHD) ELISA Kit | RHD elisa kit

Human Rh Blood Group, D Antigen (RHD) ELISA Kit

CATALOG #
AAA68078

REACTIVITY
Human


PRODUCT SYNONYMS

Rh Blood Group; D Antigen (RHD); Human Rh Blood Group; D Antigen (RHD) ELISA Kit; RHD elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of RHD. No significant cross-reactivity or interference between RHD and analogues was observed.
Sequence Length	378
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	0.058ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level RHD were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level RHD were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for RHD elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of RHD in human tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to RHD. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to RHD. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain RHD, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of RHD in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	189095266
NCBI GeneID	6007
NCBI Accession #	NP_001121163.1
NCBI GenBank Nucleotide #	NM_001127691.2
UniProt Accession #	Q02161; Q02162; Q07618; Q16147; Q16235; Q16355; Q5VSK0; Q5XLS9; Q5XLT1; Q5XLT2; Q9NPK0; Q9UQ20
Molecular Weight	41,694 Da
NCBI Official Full Name	blood group Rh(D) polypeptide isoform 2
NCBI Official Synonym Full Names	Rh blood group D antigen
NCBI Official Symbol	RHD
NCBI Official Synonym Symbols	RH; Rh4; RH30; RhII; RhPI; DIIIc; RHCED; RHDel; RHPII; RhDCw; CD240D; RHXIII; RHDVA(TT); RhK562-II
NCBI Protein Information	blood group Rh(D) polypeptide
UniProt Protein Name	Blood group Rh(D) polypeptide
UniProt Gene Name	RHD
UniProt Synonym Gene Names	RhPII

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.