

Human Sarcoglycan Delta (SGCd) ELISA Kit | SGCd elisa kit

Human Sarcoglycan Delta (SGCd) ELISA Kit

CATALOG #
AAA68079

REACTIVITY
Human


PRODUCT SYNONYMS

Sarcoglycan Delta (SGCd); Human Sarcoglycan Delta (SGCd) ELISA Kit; SGCd elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of SGCd. No significant cross-reactivity or interference between SGCd and analogues was observed.
Sequence Length	256
Samples	Tissue homogenates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.781-50ng/mL
Sensitivity	0.24ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SGCd were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SGCd were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SGCD elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of SGCD in human tissue homogenates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SGCD. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SGCD. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SGCD, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SGCD in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	21314619
NCBI GeneID	6444
NCBI Accession #	NP_000328.2
NCBI GenBank Nucleotide #	NM_000337.5
UniProt Accession #	Q92629; Q53XA5; Q99644; A8K9S9
Molecular Weight	28,750 Da
NCBI Official Full Name	delta-sarcoglycan isoform 1
NCBI Official Synonym Full Names	sarcoglycan delta
NCBI Official Symbol	SGCD
NCBI Official Synonym Symbols	SGD; DAGD; 35DAG; CMD1L; SGCDP; SG-delta
NCBI Protein Information	delta-sarcoglycan
UniProt Protein Name	Delta-sarcoglycan
UniProt Gene Name	SGCD
UniProt Synonym Gene Names	Delta-SG; 35DAG

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.