

Bovine Fibroblast Growth Factor 2, Basic (FGF2) ELISA Kit | FGF2 elisa kit

Bovine Fibroblast Growth Factor 2, Basic (FGF2) ELISA Kit

CATALOG #
AAA68220

REACTIVITY
Bovine


PRODUCT SYNONYMS

Fibroblast Growth Factor 2; Basic (FGF2); Bovine Fibroblast Growth Factor 2; Basic (FGF2) ELISA Kit; FGF2 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Bovine
Specificity	This assay has high sensitivity and excellent specificity for detection of FGF2. No significant cross-reactivity or interference between FGF2 and analogues was observed.
Sequence Length	155
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	15.625-1000pg/mL
Sensitivity	6.5pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FGF2 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FGF2 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FGF2 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of FGF2 in bovine serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FGF2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FGF2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FGF2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FGF2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	933123263
NCBI GeneID	281161
NCBI Accession #	NP_776481.2
NCBI GenBank Nucleotide #	NM_174056.4
UniProt Accession #	F1N3T9; P03969
Molecular Weight	17,266 Da
NCBI Official Full Name	fibroblast growth factor 2
NCBI Official Symbol	FGF2
NCBI Official Synonym Symbols	BFGF; FGF-2; HBGF-2
NCBI Protein Information	fibroblast growth factor 2
UniProt Protein Name	Fibroblast growth factor
UniProt Gene Name	FGF2
UniProt Synonym Gene Names	FGF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/bovine-ovine-elisa-kits/fibroblast-growth-factor-2-basic-fgf2/68220>