

Spleen Focus Forming Virus Proviral Integration Oncogene (SPI1) ELISA Kit | SPI1 elisa kit

Rat Spleen Focus Forming Virus Proviral Integration Oncogene (SPI1) ELISA Kit

CATALOG #
AAA68266

REACTIVITY
Rat


PRODUCT SYNONYMS

Spleen Focus Forming Virus Proviral Integration Oncogene (SPI1); Rat Spleen Focus Forming Virus Proviral Integration Oncogene (SPI1) ELISA Kit; SPI1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of SPI1. No significant cross-reactivity or interference between SPI1 and analogues was observed.
Sequence Length	271
Samples	Tissue homogenates, cell lysates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	0.116ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SPI1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SPI1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SPI1 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of SPI1 in rat tissue homogenates, cell lysates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SPI1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SPI1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SPI1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SPI1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	54312108
NCBI GeneID	366126
NCBI Accession #	NP_001005892.1
NCBI GenBank Nucleotide #	NM_001005892.2
UniProt Accession #	Q6BDS1; Q5BK69
Molecular Weight	31,089 Da
NCBI Official Full Name	transcription factor PU.1
NCBI Official Synonym Full Names	Spi-1 proto-oncogene
NCBI Official Symbol	Spi1
NCBI Official Synonym Symbols	Pu.1; Sfpi1
NCBI Protein Information	transcription factor PU.1
UniProt Protein Name	Transcription factor PU.1
UniProt Gene Name	Spi1
UniProt Synonym Gene Names	Sfpi1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.