

Porcine Stromal Cell Derived Factor 1 (SDF1) ELISA Kit | SDF1 elisa kit

Porcine Stromal Cell Derived Factor 1 (SDF1) ELISA Kit

CATALOG #
AAA68303

REACTIVITY
Porcine


PRODUCT SYNONYMS

Stromal Cell Derived Factor 1 (SDF1); Porcine Stromal Cell Derived Factor 1 (SDF1) ELISA Kit; SDF1 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Specificity	This assay has high sensitivity and excellent specificity for detection of SDF1. No significant cross-reactivity or interference between SDF1 and analogues was observed.
Sequence Length	116
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Assay Type	Sandwich
Detection Range	0.625-40ng/mL
Sensitivity	0.245ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level SDF1 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level SDF1 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kit, all reagents should be kept according to the labels on vials. The TMB Substrate, Wash Buffer, Stop Solution should be stored at 4 degree C. All others should be stored at -20 degree C.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SDF1 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of SDF1 in porcine serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SDF1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SDF1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SDF1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of SDF1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	57528015
NCBI GeneID	494460
NCBI Accession #	NP_001009580.1
NCBI GenBank Nucleotide #	NM_001009580.1
UniProt Accession #	Q6EKW4
Molecular Weight	12,622 Da
NCBI Official Full Name	stromal cell-derived factor 1
NCBI Official Symbol	CXCL12
NCBI Official Synonym Symbols	SDF-1
NCBI Protein Information	stromal cell-derived factor 1
UniProt Protein Name	CXCL12 chemokine
UniProt Gene Name	CXCL12

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.