

Human Glutathione S Transferase Alpha 3 (GSTa3) ELISA Kit | GSTa3 elisa kit

Human Glutathione S Transferase Alpha 3 (GSTa3) ELISA Kit

CATALOG #
AAA68566

REACTIVITY
Human


PRODUCT SYNONYMS

Glutathione S Transferase Alpha 3 (GSTa3); Human Glutathione S Transferase Alpha 3 (GSTa3) ELISA Kit; GST-A3; GSTA3-3; GTA3; GST class-alpha member 3; GSTa3 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of GSTa3. No significant cross-reactivity or interference between GSTa3 and analogues was observed.
Sequence Length	222
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	<0.053ng/mL.
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level GSTa3 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level GSTa3 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for GSTa3 elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of GSTa3 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to GSTa3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to GSTa3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain GSTa3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of GSTa3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	24430144
NCBI GeneID	2940
NCBI Accession #	NP_000838.3
NCBI GenBank Nucleotide #	NM_000847.4
UniProt Accession #	Q16772; O43468; Q068V6; Q8WWA8; Q9H415
Molecular Weight	25,302 Da
NCBI Official Full Name	glutathione S-transferase A3
NCBI Official Synonym Full Names	glutathione S-transferase alpha 3
NCBI Official Symbol	GSTA3
NCBI Official Synonym Symbols	GTA3; GSTA3-3
NCBI Protein Information	glutathione S-transferase A3
UniProt Protein Name	Glutathione S-transferase A3
UniProt Gene Name	GSTA3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.