

Rat Aromatase (ARO) RTU ELISA Kit | ARO rtu elisa kit

Rat Aromatase (ARO) ELISA Kit

CATALOG #
AAA69214

REACTIVITY
Rat


PRODUCT SYNONYMS

Aromatase (ARO); Rat Aromatase (ARO) ELISA Kit; CYP19A1; CYPXIX; ARO; ARO1; CPV1; CYAR; CYP19; P450AROM; P450-AROM; Estrogen synthase; Cytochrome P450; Family 19; Subfamily A; Polypeptide 1; ARO rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Specificity	This assay has high sensitivity and excellent specificity for detection of ARO. No significant cross-reactivity or interference between ARO and analogues was observed.
Sequence Length	508
Assay Type	Quantitative Sandwich
Samples	Rat serum, plasma, tissue homogenates or other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.061ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ARO were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ARO were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ARO rtu elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ARO. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ARO. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ARO, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of ARO in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of ARO in rat serum, plasma, tissue homogenates or other biological fluids.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	117295
NCBI GeneID	25147
UniProt Accession #	P22443
Molecular Weight	58,412 Da
NCBI Official Full Name	Aromatase
NCBI Official Synonym Full Names	cytochrome P450, family 19, subfamily a, polypeptide 1
NCBI Official Symbol	Cyp19a1
NCBI Official Synonym Symbols	Cyp19; Cyp19a; p450arom; Aromatase
NCBI Protein Information	aromatase
UniProt Protein Name	Aromatase
UniProt Gene Name	Cyp19a1
UniProt Synonym Gene Names	Arom; Cyp19

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.