

Human Activating Transcription Factor 4 (ATF4) RTU ELISA Kit | ATF4 rtu elisa kit

Human Activating Transcription Factor 4 (ATF4) ELISA Kit

CATALOG #
AAA69244

REACTIVITY
Human


PRODUCT SYNONYMS

Activating Transcription Factor 4 (ATF4); Human Activating Transcription Factor 4 (ATF4) ELISA Kit; TAXREB67; TXREB; CREB-2; DNA-binding protein TAXREB67; Cyclic AMP-responsive element-binding protein 2; Tax-responsive enhancer element-binding protein 67; ATF4 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	351
Assay Type	Sandwich
Samples	Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.057ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ATF4 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ATF4. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ATF4. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ATF4, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of ATF4 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	33469976
NCBI GeneID	468
NCBI Accession #	NP_001666.2
NCBI GenBank Nucleotide #	NM_001675.4
UniProt Accession #	P18848; Q96AQ3; Q9UH31
Molecular Weight	38,590 Da
NCBI Official Full Name	cyclic AMP-dependent transcription factor ATF-4
NCBI Official Synonym Full Names	activating transcription factor 4
NCBI Official Symbol	ATF4
NCBI Official Synonym Symbols	CREB2; TXREB; CREB-2; TAXREB67
NCBI Protein Information	cyclic AMP-dependent transcription factor ATF-4
UniProt Protein Name	Cyclic AMP-dependent transcription factor ATF-4
UniProt Gene Name	ATF4
UniProt Synonym Gene Names	CREB2; TXREB; cAMP-dependent transcription factor ATF-4; CREB-2; cAMP-responsive element-binding protein 2; TaxREB67

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/activating-transcription-factor-4-atf4/69244>