

Human ATPase, Cu⁺⁺ Transporting Beta Polypeptide (ATP7b) RTU ELISA Kit | ATP7b rtu elisa kit

Human ATPase, Cu⁺⁺ Transporting Beta Polypeptide (ATP7b) ELISA Kit

CATALOG #
AAA69276

REACTIVITY
Human


PRODUCT SYNONYMS

ATPase; Cu⁺⁺ Transporting Beta Polypeptide (ATP7b); Human ATPase; Cu⁺⁺ Transporting Beta Polypeptide (ATP7b) ELISA Kit; PWD; WC1; WD; WND; Wilson Disease Protein; Copper pump 2; Wilson disease-associated protein; ATP7b rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ATP7b. No significant cross-reactivity or interference between ATP7b and analogues was observed.
Sequence Length	1465
Samples	Human tissue homogenates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.156-10ng/mL
Sensitivity	< 0.056ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ATP7b were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ATP7b were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ATP7b rtu elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of ATP7b in human tissue homogenates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ATP7b. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ATP7b. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ATP7b, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of ATP7b in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	55743071
NCBI GeneID	540
NCBI Accession #	NP_000044.2
NCBI GenBank Nucleotide #	NM_000053.3
UniProt Accession #	P35670; Q16318; Q16319; Q4U3V3; Q59FJ9; Q5T7X7
Molecular Weight	26,021 Da
NCBI Official Full Name	copper-transporting ATPase 2 isoform a
NCBI Official Synonym Full Names	ATPase copper transporting beta
NCBI Official Symbol	ATP7B
NCBI Official Synonym Symbols	WD; PWD; WC1; WND
NCBI Protein Information	copper-transporting ATPase 2
UniProt Protein Name	Copper-transporting ATPase 2
UniProt Gene Name	ATP7B
UniProt Synonym Gene Names	PWD; WC1; WND

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.