

Mouse B-Lymphocyte Chemoattractant 1 (BLC1) RTU ELISA Kit | BLC1 rtu elisa kit

Mouse B-Lymphocyte Chemoattractant 1 (BLC1) ELISA Kit

CATALOG #
AAA69352

REACTIVITY
Mouse


PRODUCT SYNONYMS

B-Lymphocyte Chemoattractant 1 (BLC1); Mouse B-Lymphocyte Chemoattractant 1 (BLC1) ELISA Kit; CXCL13; SCYB13; BLC; BCA1; BLR1L; ANGIE; ANGIE2; Chemokine(C-X-C-Motif)ligand 13; Small Inducible Cytokine B Subfamily(Cys-X-Cys Motif)Member 13; BLC1 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	109
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	15.6-1000pg/mL
Sensitivity	5.9pg/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for BLC1 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to BLC1. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to BLC1. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain BLC1, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of BLC1 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	9256521
NCBI GeneID	55985
NCBI Accession #	NP_061354.1
NCBI GenBank Nucleotide #	NM_018866.2
UniProt Accession #	O55038
Molecular Weight	11,927 Da
NCBI Official Full Name	C-X-C motif chemokine 13
NCBI Official Synonym Full Names	chemokine (C-X-C motif) ligand 13
NCBI Official Symbol	Cxcl13
NCBI Official Synonym Symbols	BLC; Angie; BCA-1; BLR1L; ANGIE2; Scyb13; 4631412M08Rik
NCBI Protein Information	C-X-C motif chemokine 13
UniProt Protein Name	C-X-C motif chemokine 13
UniProt Gene Name	Cxcl13
UniProt Synonym Gene Names	Blc; Scyb13

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/b-lymphocyte-chemoattractant-1-blc1/69352>