

Mouse Cyclin Dependent Kinase Inhibitor 2A (CDKN2A) RTU ELISA Kit | CDKN2A rtu elisa kit

Mouse Cyclin Dependent Kinase Inhibitor 2A (CDKN2A) ELISA Kit

CATALOG # AAA69697	REACTIVITY Mouse
------------------------------	----------------------------


PRODUCT SYNONYMS

Cyclin Dependent Kinase Inhibitor 2A (CDKN2A); Mouse Cyclin Dependent Kinase Inhibitor 2A (CDKN2A) ELISA Kit; P16; ARF; CDK4I; CMM2; INK4; INK4a; MLM; MTS1; TP16; P14; P14ARF; P16INK4a; P19; Melanoma P16; Cyclin-dependent kinase 4 inhibitor A; Multiple tumor suppressor 1; CDKN2A rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	169
Assay Type	Sandwich
Samples	Tissue Homogenates or Other Biological Fluids.
Detection Range	0.312-20ng/mL
Sensitivity	0.108ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CDKN2A rtu elisa kit	Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CDKN2A. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CDKN2A. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CDKN2A, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of CDKN2A in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	6753390
NCBI GeneID	12578
NCBI Accession #	NP_034007.1
NCBI GenBank Nucleotide #	NM_009877.2
UniProt Accession #	Q64364; Q4U255; Q9QXC7; Q9R051; P51480
Molecular Weight	17,941 Da
NCBI Official Full Name	cyclin-dependent kinase inhibitor 2A p19ARF
NCBI Official Synonym Full Names	cyclin dependent kinase inhibitor 2A
NCBI Official Symbol	Cdkn2a
NCBI Official Synonym Symbols	Arf; p16; MTS1; Pctr1; p19ARF; p16INK4a; p19<ARF>; ARF-INK4a; INK4a-ARF; Ink4a/Arf; p16(INK4a)
NCBI Protein Information	cyclin-dependent kinase inhibitor 2A
UniProt Protein Name	Tumor suppressor ARF
UniProt Gene Name	Cdkn2a
UniProt Synonym Gene Names	ARF

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/cyclin-dependent-kinase-inhibitor-2a-cdkn2a/69697>