

Rat Chemokine C-X-C-Motif Ligand 16 (CXCL16) RTU ELISA Kit | CXCL16 rtu elisa kit

Rat Chemokine C-X-C-Motif Ligand 16 (CXCL16) ELISA Kit

CATALOG #
AAA70088

REACTIVITY
Rat


PRODUCT SYNONYMS

Chemokine C-X-C-Motif Ligand 16 (CXCL16); Rat Chemokine C-X-C-Motif Ligand 16 (CXCL16) ELISA Kit; SR-PSOX; CXCLG16; SRPSOX; Scavenger receptor for phosphatidylserine and oxidized low density lipoprotein; Small-inducible cytokine B16; CXCL16 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Sequence Length	247
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.058ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for CXCL16 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to CXCL16. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to CXCL16. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain CXCL16, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of CXCL16 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	62945312
NCBI GeneID	497942
NCBI Accession #	NP_001017478.1
NCBI GenBank Nucleotide #	NM_001017478.1
UniProt Accession #	Q6AXU5
Molecular Weight	27,196 Da
NCBI Official Full Name	C-X-C motif chemokine 16
NCBI Official Synonym Full Names	C-X-C motif chemokine ligand 16
NCBI Official Symbol	Cxcl16
NCBI Protein Information	C-X-C motif chemokine 16
UniProt Protein Name	C-X-C motif chemokine 16
UniProt Gene Name	Cxcl16

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-rtu-elisa-kits/chemokine-c-x-c-motif-ligand-16-cxcl16/70088>