

Porcine Factor Related Apoptosis Ligand (FASL) RTU ELISA Kit | FASL rtu elisa kit

Porcine Factor Related Apoptosis Ligand (FASL) ELISA Kit

CATALOG #
AAA70549

REACTIVITY
Porcine


PRODUCT SYNONYMS

Factor Related Apoptosis Ligand (FASL); Porcine Factor Related Apoptosis Ligand (FASL) ELISA Kit; CD178; CD95L; CD95-L; FASLG; APT1LG1; APT1-LG1; TNFSF6; Fas Antigen Ligand; TNF Superfamily Member 6; Tumor Necrosis Factor(ligand)Superfamily; Member 6; FASL rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Porcine
Sequence Length	282
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	23.4-1500pg/mL
Sensitivity	5.2pg/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FASL rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FASL. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FASL. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FASL, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of FASL in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #

[47523228](#)

NCBI GeneID	396726
NCBI Accession #	NP_998971.1
NCBI GenBank Nucleotide #	NM_213806.1
UniProt Accession #	Q9BEA8 ; Q95M04 ; Q95N10
Molecular Weight	31,756 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 6
NCBI Official Symbol	FASLG
NCBI Official Synonym Symbols	FASL; CD95-L; TNFSF6
NCBI Protein Information	tumor necrosis factor ligand superfamily member 6
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 6
UniProt Gene Name	FASLG
UniProt Synonym Gene Names	CD95L; FASL; TNFSF6; CD95-L; Fas ligand; FasL; sFasL; APL; FasL ICD; SPA

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/porcine-rtu-elisa-kits/factor-related-apoptosis-ligand-fasl/70549>