

Human Fibroblast Growth Factor 10 (FGF10) RTU ELISA Kit | FGF10 rtu elisa kit

Human Fibroblast Growth Factor 10 (FGF10) ELISA Kit

CATALOG #
AAA70611

REACTIVITY
Human


PRODUCT SYNONYMS

Fibroblast Growth Factor 10 (FGF10); Human Fibroblast Growth Factor 10 (FGF10) ELISA Kit; Keratinocyte growth factor 2; FGF10 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of FGF10. No significant cross-reactivity or interference between FGF10 and analogues was observed.
Sequence Length	208
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Detection Range	15.6-1000pg/mL
Sensitivity	< 5.7pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FGF10 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FGF10 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FGF10 rtu elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FGF10. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FGF10. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FGF10, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of FGF10 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of FGF10 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758360
NCBI GeneID	2255
NCBI Accession #	NP_004456.1
NCBI GenBank Nucleotide #	NM_004465.1
UniProt Accession #	O15520; Q6FHR3; Q6FHT6; Q96P59; C7FDY0
Molecular Weight	23,436 Da
NCBI Official Full Name	fibroblast growth factor 10
NCBI Official Synonym Full Names	fibroblast growth factor 10
NCBI Official Symbol	FGF10
NCBI Protein Information	fibroblast growth factor 10
UniProt Protein Name	Fibroblast growth factor 10
UniProt Gene Name	FGF10
UniProt Synonym Gene Names	FGF-10

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/fibroblast-growth-factor-10-fgf10/70611>