

Human Fibroblast Growth Factor 7 (FGF7) RTU ELISA Kit | FGF7 rtu elisa kit

Human Fibroblast Growth Factor 7 (FGF7) ELISA Kit

CATALOG #
AAA70661

REACTIVITY
Human


PRODUCT SYNONYMS

Fibroblast Growth Factor 7 (FGF7); Human Fibroblast Growth Factor 7 (FGF7) ELISA Kit; KGF; HBGF7; Keratinocyte Growth Factor; Heparin-binding growth factor 7; FGF7 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of FGF7. No significant cross-reactivity or interference between FGF7 and analogues was observed.
Sequence Length	194
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	15.6-1000pg/mL
Sensitivity	< 5.2pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level FGF7 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level FGF7 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for FGF7 rtu elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of FGF7 in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to FGF7. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to FGF7. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain FGF7, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of FGF7 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503705
NCBI GeneID	2252
NCBI Accession #	NP_002000.1
NCBI GenBank Nucleotide #	NM_002009.3
UniProt Accession #	P21781; Q6FGV5; Q96FG5; H0YNY5
Molecular Weight	11,520 Da
NCBI Official Full Name	fibroblast growth factor 7
NCBI Official Synonym Full Names	fibroblast growth factor 7
NCBI Official Symbol	FGF7
NCBI Official Synonym Symbols	KGF; HBGF-7
NCBI Protein Information	fibroblast growth factor 7
UniProt Protein Name	Fibroblast growth factor 7
UniProt Gene Name	FGF7
UniProt Synonym Gene Names	KGF; FGF-7; HBGF-7

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/fibroblast-growth-factor-7-fgf7/70661>