

Human Growth Regulated Oncogene Gamma (GROg) RTU ELISA Kit | GROg rtu elisa kit

Human Growth Regulated Oncogene Gamma (GROg) ELISA Kit

CATALOG #
AAA71032

REACTIVITY
Human


PRODUCT SYNONYMS

Growth Regulated Oncogene Gamma (GROg); Human Growth Regulated Oncogene Gamma (GROg) ELISA Kit; CXCL3; SCYB3; GRO3; GRO-G; MIP2-B; MIP2b; CINC2b; Chemokine(C-X-C-Motif)ligand 3; Macrophage inflammatory protein 2-beta; GROg rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	107
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	7.81-500pg/mL
Sensitivity	2.7pg/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for GROg rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to GROg. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to GROg. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain GROg, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of GROg in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	54144650
NCBI GeneID	2921
NCBI Accession #	NP_002081.2
NCBI GenBank Nucleotide #	NM_002090.2
UniProt Accession #	P19876; Q4W5H9
Molecular Weight	11,342 Da
NCBI Official Full Name	C-X-C motif chemokine 3
NCBI Official Synonym Full Names	C-X-C motif chemokine ligand 3
NCBI Official Symbol	CXCL3
NCBI Official Synonym Symbols	GRO3; GROg; MIP2B; SCYB3; MIP-2b; CINC-2b
NCBI Protein Information	C-X-C motif chemokine 3
UniProt Protein Name	C-X-C motif chemokine 3
UniProt Gene Name	CXCL3
UniProt Synonym Gene Names	GRO3; GROG; SCYB3; GRO-gamma; MIP2-beta

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/growth-regulated-oncogene-gamma-grog/71032>