

Human Guanylate Cyclase Activator 2B (GUCA2B) RTU ELISA Kit | GUCA2B rtu elisa kit

Human Guanylate Cyclase Activator 2B (GUCA2B) ELISA Kit

CATALOG #
AAA71067

REACTIVITY
Human


PRODUCT SYNONYMS

Guanylate Cyclase Activator 2B (GUCA2B); Human Guanylate Cyclase Activator 2B (GUCA2B) ELISA Kit; UGN; GCAP-II; Uroguanylin; Guanylate cyclase C-activating peptide II; GUCA2B rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of GUCA2B. No significant cross-reactivity or interference between GUCA2B and analogues was observed.
Sequence Length	112
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates or other biological fluids
Detection Range	15.6-1000pg/mL
Sensitivity	< 6.2pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level GUCA2B were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level GUCA2B were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for GUCA2B rtu elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to GUCA2B. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to GUCA2B. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain GUCA2B, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of GUCA2B in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of GUCA2B in human serum, plasma, tissue homogenates or other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	6005774
NCBI GeneID	2981
NCBI Accession #	NP_009033.1
NCBI GenBank Nucleotide #	NM_007102.2
UniProt Accession #	Q16661; Q52LV0
Molecular Weight	12,069 Da
NCBI Official Full Name	guanylate cyclase activator 2B preproprotein
NCBI Official Synonym Full Names	guanylate cyclase activator 2B
NCBI Official Symbol	GUCA2B
NCBI Official Synonym Symbols	UGN; GCAP-II
NCBI Protein Information	guanylate cyclase activator 2B
UniProt Protein Name	Guanylate cyclase activator 2B
UniProt Gene Name	GUCA2B
UniProt Synonym Gene Names	GCAP-II; UGN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/guanylate-cyclase-activator-2b-guca2b/71067>