

Mouse Histone H4 (H4) RTU ELISA Kit | H4 rtu elisa kit

Mouse Histone H4 (H4) ELISA Kit

CATALOG # AAA71086	REACTIVITY Mouse
------------------------------	----------------------------


PRODUCT SYNONYMS

Histone H4 (H4); Mouse Histone H4 (H4) ELISA Kit; H4 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of H4. No significant cross-reactivity or interference between H4 and analogues was observed.
Sequence Length	103
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates or other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.049ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level H4 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level H4 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for H4 rtu elisa kit	Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to H4. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to H4. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain H4, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of H4 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of H4 in mouse serum, plasma, tissue homogenates or other biological fluids.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	306482623
NCBI GeneID	100041230
NCBI Accession #	NP_001182350.1
NCBI GenBank Nucleotide #	NM_001195421.1
UniProt Accession #	P62806; P02304; P02305; Q0VDL9; Q2M2Q5; Q5T006; Q6PDS7; Q811M0; Q9D0C9; A0AUM5; A4FUP8; A4QMY0
Molecular Weight	11,367 Da
NCBI Official Full Name	histone H4
NCBI Official Synonym Full Names	histone cluster 1, H4m
NCBI Official Symbol	Hist1h4m
NCBI Official Synonym Symbols	Gm11275
NCBI Protein Information	histone H4
UniProt Protein Name	Histone H4
UniProt Gene Name	Hist1h4a
UniProt Synonym Gene Names	H4-53

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/histone-h4-h4/71086>