

Human Hepatoma Derived Growth Factor (HDGF) RTU ELISA Kit | HDGF rtu elisa kit

Human Hepatoma Derived Growth Factor (HDGF) ELISA Kit

CATALOG #
AAA71124

REACTIVITY
Human


PRODUCT SYNONYMS

Hepatoma Derived Growth Factor (HDGF); Human Hepatoma Derived Growth Factor (HDGF) ELISA Kit; HMG1L2; High-Mobility Group Protein 1-Like 2; HDGF rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of HDGF. No significant cross-reactivity or interference between HDGF and analogues was observed.
Sequence Length	256
Samples	Serum, plasma, tissue homogenates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	0.312-20ng/mL
Sensitivity	< 0.112ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level HDGF were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level HDGF were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HDGF rtu elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of HDGF in human serum, plasma, tissue homogenates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HDGF. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to HDGF. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HDGF, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of HDGF in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	186928819
NCBI GeneID	3068
NCBI Accession #	NP_001119522.1
NCBI GenBank Nucleotide #	NM_001126050.1
UniProt Accession #	P51858; Q5SZ07; Q5SZ08; Q5SZ09; B3KU21; D3DVC9
Molecular Weight	28,264 Da
NCBI Official Full Name	hepatoma-derived growth factor isoform b
NCBI Official Synonym Full Names	heparin binding growth factor
NCBI Official Symbol	HDGF
NCBI Official Synonym Symbols	HMG1L2
NCBI Protein Information	hepatoma-derived growth factor
UniProt Protein Name	Hepatoma-derived growth factor
UniProt Gene Name	HDGF
UniProt Synonym Gene Names	HMG1L2; HDGF; HMG-1L2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/hepatoma-derived-growth-factor-hdgf/71124>