

Human High Mobility Group AT Hook Protein 2 (HMGA2) RTU ELISA Kit | HMGA2 rtu elisa kit

Human High Mobility Group AT Hook Protein 2 (HMGA2) ELISA Kit

CATALOG # AAA71175	REACTIVITY Human
------------------------------	----------------------------


PRODUCT SYNONYMS

High Mobility Group AT Hook Protein 2 (HMGA2); Human High Mobility Group AT Hook Protein 2 (HMGA2) ELISA Kit; BABL; HMGI-C; HMGIC; LIPO; High mobility group protein HMGI-C; HMGA2 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	90
Assay Type	Sandwich
Samples	Tissue Homogenates, Cell Lysates or Other Biological Fluids.
Detection Range	0.312-20ng/mL
Sensitivity	0.114ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HMGA2 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HMGA2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to HMGA2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HMGA2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of HMGA2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	1057503228
NCBI GeneID	8091
NCBI Accession #	NP_001317119.1
NCBI GenBank Nucleotide #	NM_001330190.1
UniProt Accession #	P52926; Q1M182; Q1M185; Q1M186; Q1M187; Q1M188; E7EP85; E7EWA2
Molecular Weight	9,742 Da
NCBI Official Full Name	high mobility group protein HMGI-C isoform 3
NCBI Official Synonym Full Names	high mobility group AT-hook 2
NCBI Official Symbol	HMGA2
NCBI Official Synonym Symbols	BABL; LIPO; HMGIC; HMGI-C; STQTL9
NCBI Protein Information	high mobility group protein HMGI-C
UniProt Protein Name	High mobility group protein HMGI-C
UniProt Gene Name	HMGA2
UniProt Synonym Gene Names	HMGIC

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/high-mobility-group-at-hook-protein-2-hmga2/71175>