

Human 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) RTU ELISA Kit | HSD17b3 rtu elisa kit

Human 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) ELISA Kit

CATALOG #
AAA71227

REACTIVITY
Human


PRODUCT SYNONYMS

17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3); Human 17-Beta-Hydroxysteroid Dehydrogenase Type 3 (HSD17b3) ELISA Kit; SDR12C2; EDH17B3; Short Chain Dehydrogenase/Reductase Family 12C; Member 2; Testosterone 17-Beta-Dehydrogenase 3; Testicular 17-beta-hydroxysteroid dehydrogenase; HSD17b3 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	310
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates or Other Biological Fluids.
Detection Range	15.6-1000pg/mL
Sensitivity	6.4pg/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HSD17b3 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to HSD17b3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to HSD17b3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HSD17b3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of HSD17b3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557649
NCBI GeneID	3293
NCBI Accession #	NP_000188.1
NCBI GenBank Nucleotide #	NM_000197.1
UniProt Accession #	P37058; Q5U0Q6
Molecular Weight	29,109 Da
NCBI Official Full Name	testosterone 17-beta-dehydrogenase 3
NCBI Official Synonym Full Names	hydroxysteroid 17-beta dehydrogenase 3
NCBI Official Symbol	HSD17B3
NCBI Official Synonym Symbols	EDH17B3; SDR12C2
NCBI Protein Information	testosterone 17-beta-dehydrogenase 3
UniProt Protein Name	Testosterone 17-beta-dehydrogenase 3
UniProt Gene Name	HSD17B3
UniProt Synonym Gene Names	EDH17B3; SDR12C2; 17-beta-HSD 3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/17-beta-hydroxysteroid-dehydrogenase-type-3-hsd17b3/71227>