

Rabbit TGFB1 Polyclonal Antibody | anti-TGFB1 antibody

Rabbit anti TGFB1 (Non-Phosphospecific) Polyclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA71349	Rabbit	Human, Mouse, Rat	ELISA, Immunoprecipitation, Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

TGFB1; Polyclonal Antibody; Rabbit anti TGFB1 (Non-Phosphospecific) Polyclonal Antibody; Transforming Growth Factor beta Receptor 1 (TGFB1); anti-TGFB1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	IgG
Specificity	This antibody recognizes human TGFB1(Nonphospho) protein. It cross reacts to human, mice and rat.
Purity/Purification	The Rabbit IgG is purified by site-modified Epitope Affinity Purification.
Form/Format	This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer
Sequence Length	507
Applicable Applications for anti-TGFB1 antibody	ELISA, IP (Immunoprecipitation), WB (Western Blot), IHC (Immunohistochemistry)
Antigen Preparation	This antibody was obtained from the depletion column of the following three Non-Phosphospecific peptides (-- IYDM-T-T-SGS--; --DMTT-S-G-S-G-S-GLPL--; --EEDP-S-LDRPFI— This sequence is also identical within human, rat, mouse, chicken, bovine.
Positive Control	Kidney Tissue
Cellular Location	Cell Membrane
Preparation and Storage	The antibodies are stable for 12 months from date of receipt when stored at -20°C to -70°C. The antibodies can be stored at 2°C-8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

IMAGES



IHC (Immunohistochemistry)
IHC: Formalin-fixed, paraffinembedded human placenta tissue stained with Rabbit anti-TGFBR 1) antibody (Cat. # AAA71349) at 1:100 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.



WB (Western Blot)
WB: The cell lysate derived from Hela (lane 1) and MCF-7 (Land 2) were immunoprobed by Rabbit anti-TGFrR1 (Cat# AAA71349) at 1:1000. A major immunoreactive band is observed around -55 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-TGFBR1 antibody	Transforming growth factor beta isoforms (TGF-beta1, 2, 3) play vital roles in regulating cellular growth and differentiation, and they signal through a highly restricted subset of receptors known as TGF-betatype I receptor (TbetaR-I) and TGF-beta type II receptor (TbetaR-II). Expression of T beta R-I is more ubiquitous than that of T beta R-II. There was no staining of T beta R-II in the central nervous system, while intense T beta R-I staining was found specifically in nervous tissues. Expression of T beta R-I and T beta R-II was mostly coincident with that of their ligands, suggesting that TGF-beta s act as multiple mediators during organogenesis. TGF-beta binds with high affinity to its receptor, a transmembrane protein with a cytoplasmic serine/threonine kinase domain, to form a dynamic dimer with phosphorylation in the cytoplasmic Ser/Thr residuals which cause its kinase activity and association with another TGF beta-binding protein, the type I receptor, to signal growth inhibition and early gene responses.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	817050443
NCBI GeneID	7046
NCBI Accession #	NP_001293139.1
NCBI GenBank Nucleotide #	NM_001306210.1
UniProt Accession #	P36897; Q6IR47; Q706C0; Q706C1
Molecular Weight	55.0
NCBI Official Full Name	TGF-beta receptor type-1 isoform 3
NCBI Official Synonym Full Names	transforming growth factor beta receptor 1
NCBI Official Symbol	TGFBR1
NCBI Official Synonym Symbols	AAT5; ALK5; ESS1; LDS1; MSSE; SKR4; ALK-5; LDS1A; LDS2A; TGFR-1; ACVRLK4; tbetaR-I
NCBI Protein Information	TGF-beta receptor type-1
UniProt Protein Name	TGF-beta receptor type-1
UniProt Gene Name	TGFBR1
UniProt Synonym Gene Names	ALK5; SKR4; TGFR-1; ALK-5; ALK5; SKR4; TGF-beta receptor type I; TbetaR-I

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/tgfb1-antibody-immunoprecipitation-western-blot-immunohistochemistry-elisa/71349>