

Mouse EGFR (Tyr-1101) Monoclonal Antibody | anti-EGFR antibody

EGFR (Tyr-1101), phospho-specific

CATALOG # AAA71505	HOST Mouse	REACTIVITY Human, Rat, Mouse	APPLICATIONS Immunocytochemistry, Western Blot, ELISA
------------------------------	----------------------	--	---


PRODUCT SYNONYMS

EGFR (Tyr-1101); Monoclonal Antibody; phospho-specific; Epidermal Growth Factor Receptor; ErbB-1; anti-EGFR antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Rat, Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Number	M199
Specificity	This antibody detects a 180 kDa* protein on SDS-PAGE immunoblots of human A431 cells treated with EGF, and does not detect this band in control cells. *All molecular weights (MW) are confirmed by comparison to MW standards and to western blot mobilities of known proteins with similar MW. "Native" western blot utilizes non-reducing sample buffer (no mercaptoethanol or SDS), normal SDS-PAGE gel electrophoresis, and no methanol in transfer buffers.
Purity/Purification	Protein G Purified
Form/Format	Mouse monoclonal, protein G purified antibody is supplied in 100ul phosphate-buffered saline, 50% glycerol, 1 mg/ml BSA, and 0.05% sodium azide.
Sequence Length	1210
Applicable Applications for anti-EGFR antibody	ICC (Immunocytochemistry), WB (Western Blot), ELISA
Immunogen	Clone M199 was generated from a phospho-EGFR (Tyr-1101) synthetic peptide (coupled to KLH) corresponding to amino acid residues surrounding tyrosine 1101 of human EGFR (ErbB-1). This human EGFR sequence has high homology to rat and mouse EGFR, and is not conserved in other ErbB family members. This site is Tyr-1125 in the P00533-1 version of the human EGFR sequence.
Preparation and Storage	Store at -20°C. Stable for 1 year.

IMAGES



WB (Western Blot)



ICC (Immunocytochemistry)

ADDITIONAL INFORMATION

Related Product Information for anti-EGFR antibody	The epidermal growth factor receptor (EGFR) is a transmembrane glycoprotein with an extracellular ligand-binding domain and a cytoplasmic domain with intrinsic tyrosine kinase activity. The cytoplasmic domain has a C-terminal region with multiple autophosphorylation sites (Tyr-992, 1068, 1086, 1148, and 1173). These sites are important for downstream signaling and rapid internalization. In addition, EGFR activation leads to c Src mediated phosphorylation of Tyr-845 and Tyr-1101. The former site is required for mitogenic responses to EGFR activation, while the latter may be an SH2 binding site. Phosphorylation of EGFR on serine and threonine residues is thought to represent a mechanism for regulation of receptor kinase activity and internalization. These sites include a PKC site (Thr-654), CAMKII sites (Ser-1046, 1047, 1057, and 1142), and constitutively phosphorylated sites (Ser-967 and Ser-1002). Thus, the regulation of EGFR activity involves a complex series of phosphorylation events at multiple sites throughout the intracellular portion of the receptor.
References	• Carpenter, G. (2000) Bioessays 22:697. • Boeri Erba, E. et al. (2005) Mol. Cell. Prot. 4:1107.

NCBI AND UNIPROT INFORMATION

NCBI GI #	29725609
NCBI GeneID	1956
NCBI Accession #	NP_005219.2
NCBI GenBank Nucleotide #	NM_005228.3
UniProt Accession #	P00533; O00688; O00732; P06268; Q14225; Q68GS5; Q92795; Q9BZS2; Q9GZX1; Q9H2C9; Q9H3C9; Q9UMD7; Q147T7; Q2TTR7; Q504U8; Q59FL8; Q68GS6; Q75MF2; Q8WZA5; Q9H3C8
NCBI Official Full Name	epidermal growth factor receptor isoform a
NCBI Official Synonym Full Names	epidermal growth factor receptor
NCBI Official Symbol	EGFR
NCBI Official Synonym Symbols	ERBB; HER1; mENA; ERBB1; PIG61
NCBI Protein Information	epidermal growth factor receptor; OTTHUMP00000159661; OTTHUMP00000159662; OTTHUMP00000159663; OTTHUMP00000209210; OTTHUMP00000209211; proto-oncogene c-ErbB-1; cell growth inhibiting protein 40; cell proliferation-inducing protein 61; receptor tyrosine-protein kinase erbB-1; avian erythroblastic leukemia viral (v-erb-b) oncogene homolog
UniProt Protein Name	Epidermal growth factor receptor
UniProt Gene Name	EGFR
UniProt Synonym Gene Names	ERBB1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/egfr-tyr-1101-antibody-clone-m199-immunocytochemistry-western-blot-elisa/71505>