

anti-Human Cytokeratin 20 Antibody | anti-KRT20 antibody

Anti-Cytokeratin 20 [10F6B12A9]

CATALOG # AAA72382	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
------------------------------	----------------------------	--


PRODUCT SYNONYMS

Cytokeratin 20; Anti-Cytokeratin 20 [10F6B12A9]; CK20; CK-20; K20; KRT20; Cytokeratin-20; Keratin-20; Keratin type I cytoskeletal 20 ;Protein IT; CGMCC No 16204; Ks20.8; anti-KRT20 antibody

PRODUCT SPECIFICATIONS


Reactivity	Human
Isotype	Mouse IgG1 kappa
Clone Number	10F6B12A9
Specificity	This antibody is specific for amino acids 411-424 of human cytokeratin 20. Cytokeratins are intermediate filaments found in the intracytoplasmic cytoskeleton of epithelial tissue. There are 19 keratin polypeptides are found in human epithelium. Cytokeratin 20 plays a significant role in maintaining keratin filament organization in intestinal epithelia. When phosphorylated, it plays a role in the secretion of mucin in the small intestine.
Purity/Purification	Purified. Protein A affinity purified
Form/Format	PBS with 0.02% Proclin 300.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-KRT20 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Preparation and Storage	Store at 4 degree C for up to 3 months. For longer storage, aliquot and store at -20 degree C.

IMAGES



WB (Western Blot)

Western blot using anti-Cytokeratin 20 antibody 10F6B12A9 . Caco-2 (A) cells lysate and human colon (B) tissue lysate (35ug protein in RIPA buffer) were resolved via SDS-PAGE, and the subsequent blots were probed with the chimeric rabbit version of 10F6B12A9 at 1ug/ml before detection using an anti-rabbit secondary antibody. A primary incubation of 1 hour was used, and proteins were detected by chemiluminescence.



IF (Immunofluorescence)

Immunofluorescence staining of Caco-2 cells with anti-Cytokeratin 20 antibody 10F6B12A9 . Immunofluorescence analysis of paraformaldehyde-fixed Caco-2 cells, permeabilized with 0.15% Triton, on Shi-fix™ coverslips stained with the chimeric rabbit IgG version of 10F6B12A9 (1:100 dilution) for 1 hour followed by Alexa Fluor 488 secondary antibody (1:1000 dilution), showing cytoplasmic and some nuclear staining. The nuclear stain is DAPI (blue). Panels show, from left to right, top to bottom, DAPI, merged channels, and an isotype control. The isotype control was an unknown specificity antibody followed by staining with Alexa Fluor 488 secondary antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-KRT20 antibody	The original antibody was generated by immunizing female ICR mice with human CK20 peptide (amino acids 411-424) conjugated to KLH in equal volume of Freund's incomplete adjuvant.
Product Categories/Family for anti-KRT20 antibody	muscle ; microfilaments ; keratin ; cancer ; oncology

NCBI AND UNIPROT INFORMATION

NCBI GI #	27894337
NCBI GeneID	54474
NCBI Accession #	NP_061883.1
NCBI GenBank Nucleotide #	NM_019010.2
UniProt Accession #	P35900 ; B2R6W7
Molecular Weight	48,487 Da
NCBI Official Full Name	keratin, type I cytoskeletal 20
NCBI Official Synonym Full Names	keratin 20
NCBI Official Symbol	KRT20
NCBI Official Synonym Symbols	K20; CD20; CK20; CK-20; KRT21
NCBI Protein Information	keratin, type I cytoskeletal 20; keratin-20; protein IT; cytokeratin 20; cytokeratin-20
UniProt Protein Name	Keratin, type I cytoskeletal 20
UniProt Gene Name	KRT20
UniProt Synonym Gene Names	CK-20; K20
UniProt Entry Name	K1C20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-antibody/cytokeratin-20/72382>