

CDK1/CyclinA2 Recombinant Protein | CDK1 recombinant protein

CDK1/CyclinA2, Active

CATALOG # AAA72961	HOST Sf9 insect cells using baculovirus	APPLICATIONS Western Blot
------------------------------	---	-------------------------------------


PRODUCT SYNONYMS

CDK1/CyclinA2; Active; CDK1 recombinant protein


PRODUCT SPECIFICATIONS

Host	Sf9 insect cells using baculovirus
Form/Format	50mM Tris-HCl, pH 7.5, 150mM NaCl, 0.25mM DTT, 0.1mM EGTA, 0.1mM EDTA, 0.1mM PMSF, 25% glycerol.
Sequence Positions	Full Length
Sequence Length	1923
Applicable Applications for CDK1 recombinant protein	WB (Western Blot), Kinase Assay
Species	Human
Tag	GST tag
Dry Ice Note	The product may be shipped with dry ice, and an additional fee may be added to your shipping cost.
Preparation and Storage	Store product at -70 degree C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

IMAGES



Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for CDK1 recombinant protein	Description: Recombinant full-length human CDK1 and CyclinA2 were co-expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag on both proteins. Scientific Background: CDK1 or Cell Division Control protein 1 is essential for the completion of START, the controlling event in the cell cycle that is required to initiate mitosis. CDK1 is a catalytic subunit of a protein kinase complex, called the M-Phase Promoting Factor that induces entry into mitosis and is universal among eukaryotes (1). Phosphorylation of Bcl-2 in G2/M phase-arrested cells following photodynamic therapy with hypericin involves a CDK1-mediated signal and delays the onset of apoptosis. Therapeutic potential of CDK inhibitor NU2058 in androgen-independent prostate cancer has also been demonstrated (2).
Product Categories/Family for CDK1 recombinant protein	Enzymes; Active Kinases

References	1. Vantieghem, A. et al: Phosphorylation of Bcl-2 in G2/M phase-arrested cells following photodynamic therapy with hypericin involves a CDK1-mediated signal and delays the onset of apoptosis. J. Biol. Chem. 2002; 277 (40):37718-31. 2. Rigas, A.C. et al: Therapeutic potential of CDK inhibitor NU2058 in androgen-independent prostate cancer. Oncogene. 2007; 18.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	281427275
NCBI GeneID	983
NCBI Accession #	NM_001786.4
NCBI GenBank Nucleotide #	NM_001786
UniProt Accession #	P06493; O60764; A8K7C4; C9J497
Molecular Weight	~59 kDa; ~78 kDa
NCBI Official Full Name	Homo sapiens cyclin dependent kinase 1 (CDK1), transcript variant 1, mRNA
NCBI Official Synonym Full Names	cyclin dependent kinase 1
NCBI Official Symbol	CDK1
NCBI Official Synonym Symbols	CDC2; CDC28A; P34CDC2
NCBI Protein Information	cyclin-dependent kinase 1
UniProt Protein Name	Cyclin-dependent kinase 1
UniProt Gene Name	CDK1
UniProt Synonym Gene Names	CDC2; CDC28A; CDKN1; P34CDC2; CDK1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/cdk1-recombinant-protein-sf9-insect-cells-using-baculovirus-wb-western-blot-kinase-assay/72961>