

FLT3 (N676K) Recombinant Protein | FLT3 recombinant protein

FLT3 (N676K), Active

CATALOG # AAA73570	HOST Sf9 insect cells	APPLICATIONS Kinase Assay
------------------------------	---------------------------------	-------------------------------------


PRODUCT SYNONYMS

FLT3 (N676K); Active; FLK2; STK1; CD135; FLT3 recombinant protein

PRODUCT SPECIFICATIONS


Host	Sf9 insect cells
Form/Format	Recombinant protein stored in 50mM Tris-HCl, pH7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol.
Applicable Applications for FLT3 recombinant protein	Kinase Assay
Tag	GST
Species	Human
Dry Ice Note	The product may be shipped with dry ice, and an additional fee may be added to your shipping cost.
Preparation and Storage	Store product at -70 degree C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

IMAGES



SDS-PAGE

Purity: Sample Purity Data. For specific information on a given lot, see related technical data sheet.



Activity

Specific Activity: Sample Kinase Activity Plot. For specific information on a given lot, see related technical data sheet.

ADDITIONAL INFORMATION

Related Product Information for FLT3 recombinant protein	Recombinant human FLT3 (N676K)(571-993) was expressed by baculovirus in Sf9 insect cells using an N-terminal GST tag. The gene accession number is NM_004119. FLT3 is a receptor tyrosine kinase that has been shown to play a role in proliferation and survival of hematopoietic progenitor cells as well as differentiation of early B lymphoid progenitors (1). FLT3 consists of an extracellular domain composed of five immunoglobulin-like domains, one transmembrane region, and a cytoplasmic kinase domain split into two parts by a kinase-insert domain. FLT3 is the most frequently mutated gene in cases of acute myelogenous leukemia (AML). About 30% to 35% of patients have either internal tandem duplications (ITDs) in the juxtamembrane domain or mutations in the activating loop of FLT3 (2). The consequence of either FLT3-ITD or activating loop mutations is the constitutive activation of the tyrosine kinase activity.
Product Categories/Family for FLT3 recombinant protein	Mutant Kinases; AKT/PKB Pathway; Angiogenesis; Cancer; ERK/MAPK Pathway; Receptor Tyrosine Kinases

References

1. Christensen, J L. et al : Flk-2 is a marker in hematopoietic stem cell differentiation: a simple method to isolate long-term stem cells. Proc. Nat. Acad. Sci. 98: 14541-14546, 2001.
2. Gilliland, D G. et al: Role of FLT3 in leukemia. Curr Opin Hematol. 2002 Jul;9(4):274-81.

NCBI AND UNIPROT INFORMATION

NCBI GenBank Nucleotide #

[NM_004119](#)

Molecular Weight

~75kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/flt3-n676k-recombinant-protein-sf9-insect-cells-kinase-assay/73570>