

Goat anti Bovine IgG (biotin) Secondary Antibody | IgG secondary antibody

Goat anti Bovine IgG (biotin)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA75477	Goat	Bovine IgG	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

Goat anti Bovine IgG (biotin); Goat anti Bovine Immunoglobulin G secondary (biotin); IgG secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Bovine IgG
Form/Format	Lyophilized from 0.01M Sodium Phosphate, 0.015M Sodium Chloride, pH7.2 with 10mg/mL BSA (Immunoglobulin and Protease free) and 0.01% Sodium Azide as preservative. Reconstitute with 1.0mL deionized water or equivalent.
Concentration	2.0 mg per vial, lyophilized (varies by lot)
Applicable Applications for IgG secondary antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Crossreactivity	Immunoelectrophoresis resulted in single precipitin arc against anti-biotin, anti-Goat Serum, Bovine IgG , Bovine IgG F(ab') ₂ , and Bovine Serum. No reaction observed against Bovine IgG F(c).
Tag/Conjugate	Biotin
Immunogen	Bovine IgG F(ab') ₂ fragment.
Preparation and Storage	Upon receipt, store at 4°C while lyophilized. Dilute only prior to immediate use. Once reconstituted, aliquot and freeze at -20C or below. Avoid Repeat freeze/thaw cycles.

ADDITIONAL INFORMATION

Product Categories/Family for IgG secondary antibody	Immunology; Conjugated Secondary Antibodies
--	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/goat-bovine-ovine-secondary-antibody/goat-anti-bovine-igg-biotin/75477>