

Goat anti Monkey IgM (FITC) Secondary Antibody | IgM secondary antibody

Goat anti Monkey IgM (FITC)

CATALOG # AAA75517	HOST Goat	REACTIVITY Monkey IgM	APPLICATIONS Immunofluorescence, Flow Cytometry, Functional Assay, ELISA
------------------------------	---------------------	---------------------------------	--

PRODUCT SYNONYMS

Goat anti Monkey IgM (FITC); Goat anti Monkey Immunoglobulin M secondary (FITC); IgM secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Monkey IgM
Specificity	Monkey IgM
Purity/Purification	Goat anti Monkey IgM (FITC) was purified by delipidation, salt fractionation and ion exchange chromatography.
Form/Format	Lyophilized from 0.02M K3PO4, pH 7.2, with 0.15M NaCl, 10 mg/ml BSA, and 0.01% NaN3. Immunoglobulin and protease free.
Applicable Applications for IgM secondary antibody	IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Biological Significance	Immunoglobulin M, or IgM for short, is a basic antibody that is produced by B cells. It is the primary antibody against A and B antigens on red blood cells. IgM is by far the physically largest antibody in the human circulatory system. It is the first antibody to appear in response to initial exposure to antigen.
Biohazard Information	This product contains sodium azide as preservative. Although the amount of sodium azide is very small appropriate care must be taken when handling this product.
Tag/Conjugate	FITC
Immunogen	Goat anti-monkey IgM (FITC) was raised in goat using monkey IgM as the immunogen.
Preparation and Storage	Store at 4 degree C until reconstitution. Following reconstitution aliquot and freeze at -20 degree C for long term storage. Avoid repeated freeze/thaw cycles.

ADDITIONAL INFORMATION

Related Product Information for IgM secondary antibody	Goat anti Monkey IgM secondary antibody (FITC)
Product Categories/Family for IgM secondary antibody	Immunology ; Conjugated Secondary Antibodies

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/goat-monkey-secondary-antibody/goat-anti-monkey-igm-fitc/75517>