

Goat anti Monkey IgM (HRP) Secondary Antibody | IgM secondary antibody

Goat anti Monkey IgM (HRP)

CATALOG # AAA75533	HOST Goat	REACTIVITY Monkey IgM	APPLICATIONS Western Blot, Immunohistochemistry, ELISA
------------------------------	---------------------	---------------------------------	--

PRODUCT SYNONYMS

Goat anti Monkey IgM (HRP); Goat anti Monkey Immunoglobulin M secondary (HRP); IgM secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Monkey IgM
Purity/Purification	Antibody purified by Ion Exchange Chromatography
Form/Format	Lyophilized in 20mM Potassium Phosphate, pH 7.2, with 0.15 M Sodium Chloride. 10 mg/mL BSA (IgG and Protease free) and 0.01% Gentimicin Sulfate. Conjugated to HRP. Do not add sodium azide. Reconstitute with 1.0ml of deionized water.
Concentration	1.0 mg, Lyophilized (varies by lot)
Applicable Applications for IgM secondary antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Tag/Conjugate	HRP
Immunogen	Monkey IgM γ chain.
Cross Reactivity	Monkey. A single precipitin arc was observed against anti-Peroxidase, anti-Goat serum, Monkey IgM and Monkey serum, when assayed by Immunoelectrophoresis. No reaction was observed against Monkey IgA or Monkey IgG.
Preparation and Storage	Upon receipt store at 4°C while lyophilized. Stable for several weeks as an undiluted liquid at 4°C, or aliquot and store -20°C for longer periods.

IMAGES



Application Data

This table displays additional reactivity among various species of serum and IgA (+) indicates antibody reactivity to the corresponding target.



WB (Western Blot)

Western Blot of Peroxidase conjugated Monkey IgM μ antibody. Lane 1: Monkey IgM. Lane 2: none. Load: 100 ng per lane. none. Peroxidase Goat secondary antibody at 1:1,000 for 60 min at RT. Block: Blocking buffer for 30 min at RT. Observed size: 72 kDa for Rabbit IgG..

ADDITIONAL INFORMATION

Related Product Information for IgM secondary antibody	Goat anti Monkey IgM secondary antibody (HRP)
Product Categories/Family for IgM secondary antibody	Immunology ; Conjugated Secondary Antibodies

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/goat-monkey-secondary-antibody/goat-anti-monkey-igm-hrp/75533>