

Goat anti Monkey IgG Secondary Antibody | IgG secondary antibody

Goat anti Monkey IgG

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA75538	Goat	Monkey IgG	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

Goat anti Monkey IgG; Goat anti Monkey Immunoglobulin G secondary; IgG secondary antibody

PRODUCT SPECIFICATIONS


Host	Goat
Reactivity	Monkey IgG
Specificity	Monkey IgG
Purity/Purification	Goat anti Monkey IgG was purified by delipidation, selective precipitation and tandem molecular sieve chromatography followed by dialysis.
Form/Format	0.02M K3PO4, pH 7.2, with 0.15M NaCl, and 0.01% N2N3.
Concentration	1.0 mg/ml (varies by lot)
Applicable Applications for IgG secondary antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Biohazard Information	This product contains sodium azide as preservative. Although the amount of sodium azide is very small appropriate care must be taken when handling this product.
Immunogen	Goat anti-monkey IgG was raised in goat using monkey IgG gamma chain as the immunogen.
Preparation and Storage	Store at 4 degree C for short term storage. Aliquot and store at -20 degree C for long term storage. Avoid repeated freeze/thaw cycles.

IMAGES

Product	Gt-a-Monkey IgG	Gt-a-Monkey IgA	Gt-a-Monkey IgM
Rhesus Serum	+	+	+
Baboon Serum	+	+	+
Cynomolgus Serum	+	+	+
Rhesus IgG	+		
Human IgA		+	
Human IgM			+

Application Data

This table displays additional reactivity among various species of serum and IgA (+) indicates antibody reactivity to the corresponding target.



WB (Western Blot)

Western Blot of Monkey IgG gamma antibody. Lane 1: Monkey IgG. Lane 2: None. Load: 100 ng per lane. Monkey IgG gamma antibody at 1:1000 for overnight at 4 deg C. DyLight800 Goat secondary antibody at 1:20,000 for 30 min at RT. Block: Blocking buffer for 30 min at RT. Size: 55 kDa, 55 kDa for Monkey IgG Gamma..

ADDITIONAL INFORMATION

Related Product Information for IgG secondary antibody	Goat anti Monkey IgG secondary antibody
--	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/goat-monkey-secondary-antibody/goat-anti-monkey-igg/75538>