

Goat anti Human IgM (Fc specific) Secondary Antibody

Goat anti Human IgM (Fc specific)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA77571	Goat	HumanInter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this immuno- conjugate has not been tested in detail.	Immunoblot, Western Blot, Dot Blot, ELISA, Immunofluorescence

PRODUCT SYNONYMS

Goat anti Human IgM (Fc specific); Polyclonal Secondary Antibody; Goat anti Human IgM (Fc specific) secondary antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Human Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this immuno- conjugate has not been tested in detail.
Clonality	Polyclonal
Specificity	IgG fraction of polyclonal Goat antiSerum to Human IgM, Fc specific
Applicable Applications for Goat anti Human IgM (Fc specific) secondary antibody	IB (Immunoblot), DB (Dot Blot), ELISA, IF (Immunofluorescence)
Host Note	Purified homogenous IgM isolated from human serum. Immunization with intact (19S) and split IgM (7S). Freund's complete adjuvant is used in the first step of the immunization procedure.
Reactivity Note	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this immuno- conjugate has not been tested in detail.
Preparation and Storage	The lyophilized IgG fraction is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.

ADDITIONAL INFORMATION

Related Product Information for Goat anti Human IgM (Fc specific) secondary antibody	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with IgG, IgA and IgG/Fab or any non-Ig protein in human serum, as tested by immunoelectrophoresis and double radial immunodiffusion. As unlabelled primary or secondary reagent for indirect detection techniques, to prepare conjugates with markers of the user's own choice, to prepare an insoluble immunoaffinity adsorbent or a solid phase antibody reagent by coupling to an artificial carrier and as catching or detection antibody in non-isotopic methodology and solid phase immunochemistry. When applied in any cytochemical or histochemical procedure or solids phase coupling technique, the optimum concentration of the IgG preparation should always be established by titration.
Product Categories/Family for Goat anti Human IgM (Fc specific) secondary antibody	Secondary Antibody

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

