

Rabbit anti Sheep IgM (Fc specific) Secondary Antibody

Rabbit anti Sheep IgM (Fc specific), conjugated with Biotin

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA77590	Rabbit	Sheep	Immunoblot, Western Blot, Dot Blot, Immunohistochemistry, Immunocytochemistry, ELISA

PRODUCT SYNONYMS

Rabbit anti Sheep IgM (Fc specific); Polyclonal Secondary Antibody; conjugated with Biotin; Rabbit anti Sheep IgM (Fc specific) secondary antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Sheep
Clonality	Polyclonal
Specificity	Biotin-conjugated IgG fraction of polyclonal Rabbit antiSerum to Sheep IgM, Fc specific.
Purity/Purification	Hyperimmune antisera with strong precipitating activity are selected for purification of the IgG fraction by Protein G chromatography.
Applicable Applications for Rabbit anti Sheep IgM (Fc specific) secondary antibody	IB (Immunoblot), DB (Dot Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), ELISA
Source	Purified normal IgM isolated from pooled sheep serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Product type	Secondary Antibodies
Immunogen	Purified normal IgM isolated from pooled sheep serum. Freund's complete adjuvant is used in the first step of the immunization procedure. Biotin-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C
Reactivity Note	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail, however a strong reaction with Goat IgM has been observed.
Product	Biotin-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C.
Preparation and Storage	The lyophilized conjugate is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.

ADDITIONAL INFORMATION

Related Product Information for Rabbit anti Sheep IgM (Fc specific) secondary antibody	The reactivity of the antiserum is directed to the Fc subunit of the IgM molecule which expresses strict isotypic (class) specificity. It does not react with any non-Ig protein in sheep serum, as tested by immunoelectrophoresis and double radial immunodiffusion. In immunocytochemical and immunohistochemical staining for the detection of IgM at the cellular and subcellular level by staining of appropriately treated cell and tissue substrates; to demonstrate circulating IgM antibodies in microbiology and autoimmune diseases; to identify a specific antigen using an reference antibody of sheep origin known to be of the IgM isotype in the middle layer of the indirect test procedure; in non-isotopic assay methodology (e.g. ELISA) to measure IgM in sheep serum or other body fluids. As a second step an avidin or streptavidin conjugate of the user's choice has to be used. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal.
Product Categories/Family for Rabbit anti Sheep IgM (Fc specific) secondary antibody	Conjugated Secondary Antibody

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/rabbit-anti-sheep-igm-fc-specific-secondary-antibody-immunoblot-dot-blot-immunohistochemistry-immunocytochemistry-elisa/77590>