

## Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (heavy and light chains) Secondary Antibody

Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (heavy and light chains), conjugated with Horseradish peroxidase

| CATALOG # | HOST   | REACTIVITY                                                                                                                                                                                                                                                   | APPLICATIONS                                                                                |
|-----------|--------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|
| AAA77599  | Rabbit | <b>Mouse</b> Cross-reactivity: Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail. | <b>Immunoblot, Western Blot, Dot Blot, Immunohistochemistry, Immunocytochemistry, ELISA</b> |

### PRODUCT SYNONYMS

Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (heavy and light chains); Polyclonal Secondary Antibody; conjugated with Horseradish peroxidase; Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (heavy and light chains) secondary antibody

### PRODUCT SPECIFICATIONS

|                                                                                                                 |                                                                                                                                                                                                                                                                                                                               |
|-----------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Host                                                                                                            | Rabbit                                                                                                                                                                                                                                                                                                                        |
| Reactivity                                                                                                      | Mouse<br><br>Cross-reactivity: Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has not been tested in detail.                                                                  |
| Clonality                                                                                                       | Polyclonal                                                                                                                                                                                                                                                                                                                    |
| Specificity                                                                                                     | Horseradish peroxidase-conjugated IgG fraction of polyclonal Rabbit antiSerum to Mouse IgG, heavy and light chains                                                                                                                                                                                                            |
| Applicable Applications for Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (heavy and light chains) secondary antibody | IB (Immunoblot), DB (Dot Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), ELISA                                                                                                                                                                                                                                  |
| Conjugation                                                                                                     | HRP                                                                                                                                                                                                                                                                                                                           |
| Physical form                                                                                                   | Peroxidase-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2).                                                                                                                                                                                                    |
| Immunogen                                                                                                       | Purified normal IgG including all known subclasses isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.                                                                                                                                                      |
| Identity & Specificity                                                                                          | The reactivity of the antiserum is directed to the Fc and Fab subunits of the IgG molecule. It includes a certain degree of reactivity with other immunoglobulins via the common Fab portion. It does not react with any non-Ig protein in mouse serum, as tested by immunoelectrophoresis and double radial immunodiffusion. |
| Physicochemical characteristics                                                                                 | IgG protein concentration 7.0 mg/ml. Enzyme/IgG protein molar ratio (E/P) is approximately 1.7. No foreign proteins added.                                                                                                                                                                                                    |
| Enzyme marker                                                                                                   | Horseradish peroxidase enriched for isoenzyme C (RZ = 3.2).                                                                                                                                                                                                                                                                   |
| Conjugation procedure                                                                                           | Conjugation is carried out using a proprietary modification of the periodate method, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. No foreign protein has been added. The conjugate is lyophilized to assure stability and long shelf life.         |

|                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|--------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Intended Use</b>            | In enzyme-immunocytochemical and immunohistochemical staining for the detection of IgG, antigen or antibody, of appropriately treated cell and tissue substrates at the cellular and subcellular level. In non-isotopic assay methodology (e.g. ELISA) to identify and measure a specific IgG in mouse serum or other body fluid. In electron microscopy, since the complex between the conjugated antibody and the antigen also has electron-dense properties. This immunoconjugate is not pre-diluted. The optimum working dilution of each conjugate should be established by titration before being used. Excess labelled antibody must be avoided because it may cause high unspecific background staining and interfere with the specific signal. Working dilutions for histochemical and cytochemical use are usually between 1 :100 and 1 :500; in ELISA and comparable non-precipitating antibody-binding assays between 1:1,000 and 1 :10,000.                                                              |
| <b>Caution</b>                 | This immunoconjugate should be handled by qualified persons only and appropriate precautions should be taken in its handling and disposal, and of all associated materials. For in vitro laboratory research purposes only.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Preparation and Storage</b> | <p>The lyophilized conjugate is shipped at ambient temperature and may be stored at +4 degree C; prolonged storage at or below -20 degree C. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20 degree C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4 degree C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the immunoconjugate.</p> <p><b>Storage/ Shelf life:</b></p> <p>Lyophilized at +4 degree C = at least 10 years<br/> reconstituted at or below -20 degree C = 3-5 years<br/> reconstituted at +4 degree C = 7 days</p> |

## ADDITIONAL INFORMATION

|                                                                                                                          |                                      |
|--------------------------------------------------------------------------------------------------------------------------|--------------------------------------|
| <b>Product Categories/Family for Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (heavy and light chains) secondary antibody</b> | <b>Conjugated Secondary Antibody</b> |
|--------------------------------------------------------------------------------------------------------------------------|--------------------------------------|

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rabbit-anti-mouse-igg1-igg2a-igg2b-igg3-heavy-and-light-chains-secondary-antibody-immunoblot-dot-blot-immunohistochemistry-immunocytochemistry-elisa/77599>