

Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific) Secondary Antibody

Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA77603	Rabbit	Mouse	Immunofluorescence, Immunoblot, Western Blot, ELISA, Dot Blot

PRODUCT SYNONYMS

Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific); Polyclonal Secondary Antibody; Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific) secondary antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Specificity	Purified IgG fraction of polyclonal Rabbit antiSerum to Mouse IgG, Fc specific
Applicable Applications for Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific) secondary antibody	IF (Immunofluorescence), IB (Immunoblot), ELISA, DB (Dot Blot)
Host Note	Purified normal IgG, including all known subclasses, isolated from pooled mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure.
Form	Purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. No foreign protein added. It is reconstituted by adding 1 ml sterile distilled water, spun down to remove insoluble particles, divided into small aliquots, frozen and stored at or below -20°C.
Reactivity Note	Inter-species cross-reactivity is a normal feature of antibodies to immunoglobulins, since Ig of different species frequently share antigenic determinants. Cross-reactivity of this antiSerum has been tested in double radial immunodiffusion anti several sera with the following results: Bovine - Cat - Chicken - Dog - Duck - Goat - Guinea Pig+ Human - Hamster + Horse - Monkey - pigeon - Rat + Sheep - Swine - Turkey - The absence of a reaction in double radial immunodiffusion does not exclude some reaction in more sensitive techniques.
Preparation and Storage	The lyophilized IgG fraction is shipped at ambient temperature and may be stored at +4°C; prolonged storage at or below -20°C. Prior to use, an aliquot is thawed slowly at ambient temperature, spun down again and used to prepare working dilutions by adding sterile phosphate buffered saline (PBS, pH 7.2). Repeated thawing and freezing should be avoided. Working dilutions should be stored at +4°C, not refrozen, and preferably used the same day. If a slight precipitation occurs upon storage, this should be removed by centrifugation. It will not affect the performance of the product.

ADDITIONAL INFORMATION

Related Product Information for Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific) secondary antibody	The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule. The antiserum contains antibodies to subclass-specific determinants as well as to determinants shared by two or more subclasses of IgG. In immunoelectrophoresis and double radial immunodiffusion, using various anti- serum concentrations against normal mouse plasma, serum, and immunoglobulin fractions, no reaction is obtained with IgA, IgM and IgG/Fab fragments or any other serum protein. As unlabelled primary or secondary reagent for indirect detection techniques, to prepare conjugates with markers of the user's own choice, to prepare an insoluble immunoaffinity adsorbent or a solid phase antibody reagent by coupling to an artificial carrier and as catching or detection antibody in non- isotopic methodology and solid phase immunochemistry. When applied in any cytochemical or histochemical procedure or solids phase coupling technique, the optimum concentration of the IgG preparation should always be established by titration.
--	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rabbit-anti-mouse-igg1-igg2a-igg2b-igg3-fc-specific-secondary-antibody-immunofluorescence-immunoblot-dot-blot-elisa/77603>