

## Mouse anti-Chicken TCR alpha beta (V beta 1) Monoclonal Antibody

Mouse Anti-Chicken TCRab/Vb1-UNLB

| CATALOG # | HOST  | REACTIVITY | APPLICATIONS                                                                |
|-----------|-------|------------|-----------------------------------------------------------------------------|
| AAA78704  | Mouse | Chicken    | Immunoprecipitation, Immunohistochemistry, Flow Cytometry, Functional Assay |

### PRODUCT SYNONYMS

TCR alpha beta (V beta 1); Monoclonal Antibody; Mouse Anti-Chicken TCRab/Vb1-UNLB; Mouse Anti-Chicken TCRab/Vb1 - Purified; Unlabeled Antibody; Mouse Anti-Chicken TCR alpha beta (V beta 1); anti-TCR alpha beta (V beta 1) antibody

### PRODUCT SPECIFICATIONS

|                                                                     |                                                                                                                                                                                                                                              |
|---------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Host                                                                | Mouse                                                                                                                                                                                                                                        |
| Reactivity                                                          | Chicken                                                                                                                                                                                                                                      |
| Clonality                                                           | Monoclonal                                                                                                                                                                                                                                   |
| Isotype                                                             | Mouse IgG1 kappa                                                                                                                                                                                                                             |
| Clone Number                                                        | TCR-2                                                                                                                                                                                                                                        |
| Specificity                                                         | Chicken/Turkey/Peacock/Guinea Fowl TCRalpha beta/V beta <sub>1</sub> ; Mr 50& 40 kDa                                                                                                                                                         |
| Form/Format                                                         | The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added.                                                   |
| Applicable Applications for anti-TCR alpha beta (V beta 1) antibody | IP (Immunoprecipitation), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry)                                                                                                                                                              |
| Characterization                                                    | To ensure lot-to-lot consistency, each batch of monoclonal antibody is tested to conform with characteristics of a standard reference reagent using immunofluorescence staining and analysis on a Becton Dickinson FACSscan™ flow cytometer. |
| Preparation and Storage                                             | Store at 2-°C.                                                                                                                                                                                                                               |

### ADDITIONAL INFORMATION

|                                                                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
|-------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Related Product Information for anti-TCR alpha beta (V beta 1) antibody | <p>MAB TCR2 precipitates a CD3-associated heterodimer of Mr 90-kDa (two bands of Mr 40-kDa and 50- kDa upon reduction) on chicken peripheral blood T cells.<sup>1</sup> Deglycosylation of the heterodimer yields two polypeptides of Mr 34-kDa and 29-kDa from TCR2 precipitates. In the chicken, two distinct subpopulations of alpha beta T cells appear in the thymus subsequent to the appearance of gamma delta T cells. These subpopulations, originally denoted as TCR2 and TCR3,1-3 arise sequentially in the thymus during ontogeny and are now known to represent two distinct V beta families, V beta 1 and V beta 2, respectively.<sup>4</sup> The TCR2 MAB reacts with approximately 40% of thymocytes, 40-50% of blood mononuclear cells and 40% of splenocytes in the chicken. Two-color immunofluorescence has revealed that most of the TCR2+ thymocytes express both CD4 and CD8 antigens. The TCR2+ cells in blood were found to express either CD4 (74+/-2%) or CD8 (26+/-4%). TCR2+ cells in the spleen also express either CD4 (37+/-1%) or CD8 (64+/-4%). Surprisingly, a relatively large subpopulation of CD8+ cells in the spleen are negative for TCR2.1-3 This observation led to the demonstration that 71+/-6% of the TCR1(gamma delta)+ cells in the spleen express the CD8 marker.</p> |
|-------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

## References

1. Chen, C.H., J. Cihak, U. Losch, and M.D. Cooper. 1988. Eur. J. Immunol. 18:539.
2. Chen, C.H., J.T. Sowder, J.M. Lahti, J. Cihak, U. Losch, and M.D. Cooper. 1989. Proc. Natl. Acad. Sci. (USA):86:2351
3. Char, D., P. Sanchez, C.H. Chen, R.P. Bucy, and M.D. Cooper. 1990. J. Immunol. 145:3547.
4. Chen, C.H., T.W.F. Gobel, T. Kubota, and M.D. Cooper. 1993. 1994. Poultry Science 73:1012.
5. Sowder, J.T., C.H. Chen, L.L. Ager, J. Cihak, U. Losch, and M.D. Cooper. 1988b. Fed. Proc. 2:874.
6. Cihak, J., U. Losch, G. Hoffman-Hezer, C.H. CHen, M.D. Cooper, and H.W.L. Ziegler-Heitbrock. 1993. Scand. J. Immunol. 38:123.
7. Chen, C.H. 1996. Personal communication.

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | [contact@aaabiotech.com](mailto:contact@aaabiotech.com)

Online datasheet: <https://stage.aaabiotech.com/monoclonal/tcr-alpha-beta-v-beta-1-antibody-clone-tcr-2-immunoprecipitation-immunohistochemistry-flow-cytometry-elisa/78704>